

EMS ExDS Run Log

Created by Dr. Thomas R. Scaggs, MD
Carle Arrow Ambulance Medical Director

Sedation for the Extremely Agitated Patient EMS form

Run Number:

Arrow Ambulance PRO Ambulance

Before Ketamine

- +4 Excited Delirium
- +3 Extremely Agitated
- +2 Moderately Agitated
- +1 Restless
- 0 Alert, Oriented, Calm
- 1 Drowsy
- 2 Light Sedation
- 3 Moderate Sedation
- 4 Deep Sedation
- 5 Unresponsive

After Ketamine

- +4 Excited Delirium
- +3 Extremely Agitated
- +2 Moderately Agitated
- +1 Restless
- 0 Alert, Oriented, Calm
- 1 Drowsy
- 2 Light Sedation
- 3 Moderate Sedation
- 4 Deep Sedation
- 5 Unresponsive

+ 4 Excited Delirium

1. Extreme combative, violent, aggressive, agitated behavior, with amazing physical strength, struggle despite futility, unlimited endurance, pain insensitivity, constant yelling, aggression toward inanimate objects, running for no reason;
2. Delusional, paranoid, irrational, bizarre behavior, inability to maintain attention, disorientation, disorganized thinking, incoherent speech, naked/semi-clothed

+3 Extremely Agitated

1. Severe combative, violent, aggressive, agitated behavior, significant physical strength, struggle despite futility, significant endurance, pain insensitivity, constant yelling/screaming;
2. Displaying no signs of delirium

+2 Moderately Agitated

- Moderate violent/agitated behavior, frequent yelling

+1 Restless

- Anxious or apprehensive but movements are not aggressive or vigorous

0 Alert, Oriented, Calm

-1 Drowsy

- Not fully alert, but has sustained (more than 10 seconds) awakening, with eye contact to voice

-2 Light Sedation

- Briefly (less than 10 seconds) awakens with eye contact to voice

-3 Moderate Sedation

- Any movement (but no eye contact) to voice

-4 Deep Sedation

- No response to voice, any movement to physical stimulation

-5 Unresponsive

- No response to voice or physical stimulation

Medic Safety Scale:

During the restraining process what would you classify your safety?

Make a selection ▼

During the drug administration, what would you classify your safety?

Make a selection ▼

Were there any known potential causes of the excited delirium?

Yes No

Drug use?

Yes No

If so what?

Alcohol withdrawal

Yes No

Psychiatric problem:

Yes No

If so what?

Psychiatric Medication changes:

Yes No

If so what?

Other - Explain:

Was there trauma to the patient?

Yes No

If so, explain:

Were there any actions taken by police/bystanders prior to arrival?

Yes No

Did they attempt to restrain the patient?

Yes No

If so, explain:

Were there any adverse affects due to ketamine?

Yes No

Airway Maintenance

Vomiting

Oral secretions

Increased muscle tone

Emergence reaction

Other

Explain other:

What was the condition of the pupils before administration?

- Right Pupil Dilated Reactive
 Constricted Nonreactive
 Normal

- Left Pupil Dilated Reactive
 Constricted Nonreactive
 Normal

What was the time taken to restrain the patient once hands were placed on the patient?

Were there any injuries to the patient during the restraining process beside minor contusions and abrasions?

- Yes No

Explain

How was the medication administered?

- Clean stick Dirty stick

Mark behaviors the patient possessed prior to administration of medication.

- Agitation
 Violent
 Aggression toward objects
 Superhuman Strength
 Unlimited endurance
 Screaming
 Talks to invisible people
 Decreased pain sensation
 Naked or partially dressed
 Paranoia
 Disorientation
 Hallucination
 Delusional
 Easily distracted
 Running for no reason

Submit